

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 38/00</b>	<b>A1</b>	(11) International Publication Number: <b>WO 95/09000</b> (43) International Publication Date: <b>6 April 1995 (06.04.95)</b>
(21) International Application Number: <b>PCT/US94/10872</b> (22) International Filing Date: <b>26 September 1994 (26.09.94)</b> (30) Priority Data: <b>130,032 30 September 1993 (30.09.93) US</b> (60) Parent Application or Grant (63) Related by Continuation <b>US 130,032 (CON)</b> <b>Filed on 30 September 1993 (30.09.93)</b> (71) Applicant (for all designated States except US): <b>MERCK &amp; CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</b> (72) Inventors; and (75) Inventors/Applicants (for US only): <b>DESOLMS, S., Jane [US/US]; 735 Port Indian Road, Norristown, PA 19403 (US). GIULIANI, Elizabeth, A. [US/US]; 1333 Gwynedale Way, Lansdale, PA 19446 (US). GRAHAM, Samuel, L. [US/US]; 325 Widlund Drive, Schwenksville, PA 19473 (US).</b>		(74) Common Representative: <b>MERCK &amp; CO., INC.; Patent Dept., 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</b> (81) Designated States: <b>AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MN, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).</b>  <b>Published</b> <i>With international search report.</i>
(54) Title: <b>INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE</b> (57) Abstract <p>The present invention comprises analogs of the CAAX motif of the protein Ras that is modified by farnesylation <i>in vivo</i>. These CAAX analogs inhibit the farnesylation of Ras. Furthermore, these CAAX analogs differ from those previously described as inhibitors of Ras farnesyl transferase in that they do not have a thiol moiety. The lack of the thiol offers unique advantages in terms of improved pharmacokinetic behavior in animals, prevention of thiol-dependent chemical reactions, such as rapid autooxidation and disulfide formation with endogenous thiols, and reduced systemic toxicity. Further contained in this invention are chemotherapeutic compositions containing these farnesyl transferase inhibitors and methods for their production.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

TITLE OF THE INVENTION

## INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE

BACKGROUND OF THE INVENTION

5           The Ras gene is found activated in many human cancers, including colorectal carcinoma, exocrine pancreatic carcinoma, and myeloid leukemias. Biological and biochemical studies of Ras action indicate that Ras functions like a G-regulatory protein, since Ras must be localized in the plasma membrane and must bind with GTP in order  
10       to transform cells (Gibbs, J. et al., *Microbiol. Rev.* **53**:171-286(1989). Forms of Ras in cancer cells have mutations that distinguish the protein from Ras in normal cells.

          At least 3 post-translational modifications are involved with Ras membrane localization, and all 3 modifications occur at the C-  
15       terminus of Ras. The Ras C-terminus contains a sequence motif termed a "CAAX" or "Cys-Aaa1-Aaa2-Xaa" box (Aaa is an aliphatic amino acid, the Xaa is any amino acid) (Willumsen et al., *Nature* **310**:583-586 (1984)). Other proteins having this motif include the Ras-related GTP-binding proteins such as Rho, fungal mating factors, the nuclear lamins,  
20       and the gamma subunit of transducin.

          Farnesylation of Ras by the isoprenoid farnesyl pyrophosphate (FPP) occurs *in vivo* on Cys to form a thioether linkage (Hancock et al., *Cell* **57**:1167 (1989); Casey et al, *Proc. Natl Acad. Sci. USA* **86**:8323 (1989)). In addition, Ha-Ras and N-Ras are  
25       palmitoylated *via* formation of a thioester on a Cys residue near a C-terminal Cys farnesyl acceptor (Gutierrez et al., *EMBO J* **8**:1093-1098 (1989); Hancock et al., *Cell* **57**: 1167-1177 (1989); Ki-Ras lacks the palmitate acceptor Cys. The last 3 amino acids at the Ras C-terminal end are removed proteolytically, and methyl esterification occurs at the  
30       new C-terminus (Hancock et al, *ibid*). Fungal mating factor and mammalian nuclear lamins undergo identical modification steps (Anderegg et al, *J. Biol. Chem.* **263**:18236 (1988); Farnsworth et al., *J. Biol. Chem.* **264**:20422 (1989)).

- 2 -

Inhibition of Ras farnesylation in vivo has been demonstrated with lovastatin (Merck & Co., Rahway, NJ) and compactin (Hancock et al., ibid; Casey et al., ibid; Schafer et al., Science 245:379 (1989)). These drugs inhibit HMG-CoA reductase, the rate limiting enzyme for the production of polyisoprenoids and the farnesyl pyrophosphate precursor. It has been shown that a farnesyl-protein transferase using farnesyl pyrophosphate as a precursor is responsible for Ras farnesylation. (Reiss et al., Cell, 62: 81-88 (1990); Schaber et al., J. Biol. Chem., 265:14701-14704 (1990); Schafer et al., Science, 249: 1133-1139 (1990); Manne et al., Proc Natl. Acad. Sci USA, 87: 7541-7545 (1990)).

Inhibition of farnesyl-protein transferase and, thereby, of farnesylation of the Ras protein, blocks the ability of Ras to transform normal cells to cancer cells. The compounds of the invention inhibit Ras farnesylation and, thereby, generate soluble Ras which, as indicated infra, can act as a dominant negative inhibitor of Ras function. While soluble Ras in cancer cells can become a dominant negative inhibitor, soluble Ras in normal cells would not be an inhibitor.

A cytosol-localized (no Cys-Aaa1-Aaa2-Xaa box farnesylation signal domain present) and activated (impaired GTPase activity, staying bound to GTP) form of Ras acts as a dominant negative Ras inhibitor of membrane-bound Ras function (Gibbs et al., Proc. Natl. Acad. Sci. USA 86:6630-6634(1989)). Cytosol-localized forms of Ras with normal GTPase activity do not act as inhibitors. Gibbs et al., ibid, showed this effect in Xenopus oocytes and in mammalian cells.

Administration of compounds of the invention to block Ras farnesylation not only decreases the amount of Ras in the membrane but also generates a cytosolic pool of Ras. In tumor cells having activated Ras, the cytosolic pool acts as another antagonist of membrane-bound Ras function. In normal cells having normal Ras, the cytosolic pool of Ras does not act as an antagonist. In the absence of complete inhibition of farnesylation, other farnesylated proteins are able to continue with their functions.

- 3 -

Farnesyl-protein transferase activity may be reduced or completely inhibited by adjusting the compound dose. Partial reduction of farnesyl-protein transferase enzyme activity by adjusting the compound dose would be useful for avoiding possible undesirable side effects resulting from interference with other metabolic processes which utilize the enzyme.

These compounds and their analogs are inhibitors of farnesyl-protein transferase. Farnesyl-protein transferase utilizes farnesyl pyrophosphate to covalently modify the Cys thiol group of the Ras Cys-Aaa<sup>1</sup>-Aaa<sup>2</sup>-Xaa (CAAX) box with a farnesyl group. While inhibition of farnesyl pyrophosphate biosynthesis by inhibiting HMG-CoA reductase blocks Ras membrane localization *in vivo* and inhibits Ras function, inhibition of farnesyl-protein transferase is more specific and is attended by fewer side effects than is the case for a general inhibitor of isoprene biosynthesis.

Previously, it has been demonstrated that tetrapeptides containing cysteine as an amino terminal residue in a CAAX sequence inhibit Ras farnesylation (Schaber *et al.*, *ibid*; Reiss *et al.*, *ibid*; Reiss *et al.*, *PNAS*, 88:732-736 (1991)). Such inhibitors may inhibit while serving as alternate substrates for the Ras farnesyl-transferase enzyme, or may be purely competitive inhibitors (U.S. Patent 5,141,851, University of Texas).

Inhibitors of Ras farnesyl-protein transferase (FPTase) have been described in two general classes. The first are analogs of farnesyl diphosphate (FPP), while the second class of inhibitors is related to the protein substrate for the enzyme, Ras. Almost all of the peptide derived inhibitors that have been described are cysteine containing molecules that are related to the CAAX motif that is the signal for protein prenylation. The exception to this generalization is a class of natural products known as the peptidocinnamins (Omura, *et al.*, *J. Antibiotics* 46:222 (1993)). In general, deletion of the thiol from a CAAX derivative dramatically reduces the inhibitory potency of these compounds. However, the thiol group potentially places limitations on the therapeutic application of FPTase inhibitors with respect to

- 4 -

pharmacokinetics, pharmacodynamics and toxicity. Therefore, a functional replacement for the thiol is desirable. With the exception of the peptidocinnamins, non-thiol FPTase inhibitors that are competitive with the Ras substrate have not been described and are the subject of this invention.


It is, therefore, an object of this invention to develop tetrapeptide-based compounds which do not have a thiol moiety, and which will inhibit farnesyl transferase and the post-translational functionalization of the oncogene Ras protein. It is a further object of this invention to develop chemotherapeutic compositions containing the compounds of this invention and methods for producing the compounds of this invention.

#### SUMMARY OF THE INVENTION

The present invention comprises analogs of the CAAX motif of the protein Ras that is modified by farnesylation *in vivo*. These CAAX analogs inhibit the farnesylation of Ras. Furthermore, these CAAX analogues differ from those previously described as inhibitors of Ras farnesyl transferase in that they do not have a thiol moiety. The lack of the thiol offers unique advantages in terms of improved pharmacokinetic behavior in animals, prevention of thiol-dependent chemical reactions, such as rapid autoxidation and disulfide formation with endogenous thiols, and reduced systemic toxicity. Further contained in this invention are chemotherapeutic compositions containing these farnesyl transferase inhibitors and methods for their production.


The compounds of this invention are illustrated by the formulae:

- 5 -




5

I



10


II



20

III

and




30

IV



5

10



# I


15

30


R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;

- 7 -


X-Y is




5





10



15



20



R<sup>7</sup> is  
25

an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

Z is  
30

H<sub>2</sub> or O;

m is

0, 1 or 2; and


o is

0, 1, 2 or 3;

- 8 -

or the pharmaceutically acceptable salts thereof.

In a second embodiment of this invention the prodrugs of compounds of formula I are illustrated by the formula II:



10

II

wherein:

V is CH<sub>2</sub>, O, S, HN, or R<sup>7</sup>N;

15

20 R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;


25

30


- 9 -

X-Y is


5






10



15



20



25

R6 is

a substituted or unsubstituted aliphatic, aromatic or heteroaromatic group such as saturated chains of 1 to 8 carbon atoms, which may be branched or unbranched, wherein the aliphatic substituent may be substituted with an aromatic or heteroaromatic ring;

30

R7 is

an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

- 10 -


Z is H<sub>2</sub> or O; and

m is 0, 1 or 2;

o is 0, 1, 2 or 3;

5 or the pharmaceutically acceptable salts.

In a third embodiment of this invention, the inhibitors of farnesyl transferase are illustrated by the formula III:



15

III

wherein:

V is CH<sub>2</sub>, O, S, HN, or R<sup>7</sup>N;

20







R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;

25

30

- 11 -

X-Y is



$R^7$  is an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

Z is  $\text{H}_2$  or O;

n is 0, 1 or 2;


m is 0, 1 or 2; and

o is 0, 1, 2 or 3;

or the pharmaceutically acceptable salts thereof.

- 12 -

In a fourth embodiment of this invention the prodrugs of compounds of formula III are illustrated by the formula IV:



IV

wherein:

V is CH<sub>2</sub>, O, S, HN, or R<sup>7</sup>N;

15

20







25

30

R<sup>2</sup> and R<sup>3</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;

- 13 -

X-Y is



R<sup>7</sup> is an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

Z is H<sub>2</sub> or O;

n is 0, 1 or 2;  
 m is 0, 1 or 2; and  
 o is 0, 1, 2 or 3;  
 or the pharmaceutically acceptable salts thereof.



- 14 -

The preferred compounds of this invention are as follows:  
 N-[2(S)-(pyrrolidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine;  
 or

5 N-[2(S)-(piperidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine;  
 or the pharmaceutically acceptable salts thereof.

In the present invention, the amino acids which are  
 disclosed are identified both by conventional 3 letter and single letter  
 10 abbreviations as indicated below:

---

	Alanine	Ala	A
	Arginine	Arg	R
	Asparagine	Asn	N
15	Aspartic acid	Asp	D
	Asparagine or		
	Aspartic acid	Asx	B
	Cysteine	Cys	C
	Glutamine	Gln	Q
20	Glutamic acid	Glu	E
	Glutamine or		
	Glutamic acid	Glx	Z
	Glycine	Gly	G
	Histidine	His	H
25	Isoleucine	Ile	I
	Leucine	Leu	L
	Lysine	Lys	K
	Methionine	Met	M
	Phenylalanine	Phe	F
30	Proline	Pro	P

- 15 -

	Serine	Ser	S
	Threonine	Thr	T
	Tryptophan	Trp	W
	Tyrosine	Tyr	Y
5	Valine	Val	V

---

The compounds of the present invention may have asymmetric centers and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers, including optical isomers, being included in the present invention.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms.

The term heterocycle or heterocyclic, as used herein, represents a stable 5- to 7-membered monocyclic or stable 8- to 11-membered bicyclic or stable 11-15 membered tricyclic heterocyclic ring which is either saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O, and S, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic elements include, but are not limited to, azepinyl, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, cinnoliny, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzothiopyranyl, dihydrobenzothiopyranyl sulfone, furyl, imidazolidinyl, imidazolinyl, imidazolyl, indolinyl, indolyl, isochromanyl, isoindolinyl, isoquinolinyl, isothiazolidinyl, isothiazolyl, isothiazolidinyl, morpholinyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, 2-oxopiperazinyl, 2-oxopiperdinyl, 2-oxopyrrolidinyl, piperidyl, piperazinyl, pyridyl, pyrazinyl, pyrazolidinyl, pyrazolyl, pyrimidinyl, pyrrolidinyl, pyrrolyl,

- 16 -

quinazolinyl, quinolinyl, quinoxalinyl, tetrahydrofuryl,  
tetrahydroisoquinolinyl, tetrahydroquinolinyl, thiamorpholinyl,  
thiamorpholinyl sulfoxide, thiazolyl, thiazolinyl, thienofuryl,  
thienothienyl, and thienyl.

5           The pharmaceutically acceptable salts of the compounds of  
this invention include the conventional non-toxic salts of the compounds  
of this invention as formed, e.g., from non-toxic inorganic or organic  
acids. For example, such conventional non-toxic salts include those  
10       derived from inorganic acids such as hydrochloric, hydrobromic,  
sulfuric, sulfamic, phosphoric, nitric and the like: and the salts prepared  
from organic acids such as acetic, propionic, succinic, glycolic, stearic,  
lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic,  
phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic,  
15       fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic,  
isethionic, trifluoroacetic and the like.

          The pharmaceutically acceptable salts of the compounds of  
this invention can be synthesized from the compounds of this invention  
which contain a basic moiety by conventional chemical methods.  
20       Generally, the salts are prepared by reacting the free base with  
stoichiometric amounts or with an excess of the desired salt-forming  
inorganic or organic acid in a suitable solvent or various combinations  
of solvents.

          The compounds of the invention can be synthesized from  
their constituent amino acids by conventional peptide synthesis  
25       techniques, and the additional methods described below. Standard  
methods of peptide synthesis are disclosed, for example, in the  
following works: Schroeder *et al.*, "The Peptides", Vol. I, Academic  
Press 1965, or Bodanszky *et al.*, "Peptide Synthesis", Interscience  
Publishers, 1966, or McOmie (ed.) "Protective Groups in Organic  
30       Chemistry", Plenum Press, 1973, or Barany *et al.*, "The Peptides:  
Analysis, Synthesis, Biology" 2, Chapter 1, Academic Press, 1980, or  
Stewart *et al.*, "Solid Phase Peptide Synthesis", Second Edition, Pierce  
Chemical Company, 1984. The teachings of these works are hereby  
incorporated by reference.

- 17 -

Abbreviations used in the description of the chemistry and in the examples that follow are:

5	Ac <sub>2</sub> O	Acetic anhydride;
	Boc	t-Butoxycarbonyl;
	DBU	1,8-diazabicyclo[5.4.0]undec-7-ene;
	DMAP	4-Dimethylaminopyridine;
	DME	1,2-Dimethoxyethane;
	DMF	Dimethylformamide;
10	EDC	1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride;
	HOBT	1-Hydroxybenzotriazole hydrate;
	Et <sub>3</sub> N	Triethylamine;
	EtOAc	Ethyl acetate.
15	FAB	Fast atom bombardment;
	HOBT	3-Hydroxy-1,2,2-benzotriazin-4(3 <i>H</i> )-one;
	HPLC	High-performance liquid chromatography;
20	MCPBA	m-Chloroperoxybenzoic acid;
	MsCl	Methanesulfonyl chloride;
	NaHMDS	Sodium bis(trimethylsilyl)amide
	Py	Pyridine;
	TFA	Trifluoroacetic acid;
25	THF	Tetrahydrofuran;

Compounds of this invention are prepared by employing the reactions shown in the following Reaction Schemes A-J, in addition to other standard manipulations such as ester hydrolysis, cleavage of protecting groups, etc., as may be known in the literature or exemplified in the experimental procedures. Some key bond-forming and peptide modifying reactions are:

Reaction A. Amide bond formation and protecting group cleavage using standard solution or solid phase methodologies.

- 18 -

Reaction B. Preparation of a reduced peptide subunit by reductive alkylation of an amine by an aldehyde using sodium cyanoborohydride or other reducing agents.

5      Reaction C. Alkylation of a reduced peptide subunit with an alkyl or aralkyl halide or, alternatively, reductive alkylation of a reduced peptide subunit with an aldehyde using sodium cyanoborohydride or other reducing agents.

10      Reaction D. Peptide bond formation and protecting group cleavage using standard solution or solid phase methodologies.



15      Reaction E. Preparation of a reduced subunit by borane reduction of the amide moiety.

20      These reactions may be employed in a linear sequence to provide the compounds of the invention or they may be used to synthesize fragments which are subsequently joined by the alkylation reactions described in the Reaction Schemes.

25

30


- 19 -

REACTION SCHEME AReaction A. Coupling of residues to form an amide bondREACTION SCHEME BReaction B. Preparation of reduced peptide subunits by reductive alkylation


- 20 -

REACTION SCHEME C


Reaction C. Alkylation/reductive alkylation of reduced peptide subunits

REACTION SCHEME D

Reaction D. Coupling of residues to form an amide bond



- 21 -

REACTION SCHEME EReaction E. Preparation of reduced dipeptides from peptides

15 where  $R^A$  and  $R^B$  are  $R^2$ ,  $R^3$  or  $R^4$  as previously defined;  $X^L$  is a leaving group, e.g.,  $Br^-$ ,  $I^-$  or  $MsO^-$ ; and  $R^8$  is defined such that  $R^7$  is generated by the reductive alkylation process.

Certain compounds of this invention wherein X-Y is an ethenylene or ethylene unit are prepared by employing the reaction sequences shown in Reaction Schemes F and G. Reaction Scheme F outlines the preparation of the alkene isosteres utilizing standard manipulations such as Weinreb amide formation, Grignard reaction, acetylation, ozonolysis, Wittig reaction, ester hydrolysis, peptide coupling reaction, mesylation, cleavage of peptide protecting groups, reductive alkylation, etc., as may be known in the literature or exemplified in the Experimental Procedure. The key reactions are: stereoselective reduction of the Boc-amino-enone to the corresponding syn amino-alcohol (Scheme F, Step B, Part 1), and stereospecific boron trifluoride or zinc chloride activated organo-magnesium, organo-lithio, or organo-zinc copper(I) cyanide  $SN_2'$  displacement reaction (Scheme F, Step G). Through the use of optically pure N-Boc amino acids as starting material and these two key reactions, the stereochemistry of the final products is well defined. In Step H of Scheme F, the lactam ring is incorporated by acylation of the primary amino group with 4-chlorobutyryl chloride and base-catalyzed cyclization.




- 22 -

The alkane analogs are prepared in a similar manner by including an additional catalytic hydrogenation step as outlined in Reaction Scheme G.

5


REACTION SCHEME F

10



Step A

15




Step B

20




Step C

25


Step D  
1. LiOHStep E  
2. EDC, HOBT  
amino acid (ester)Step F  
MsCl, py

30

- 23 -

REACTION SCHEME F CONT


- 24 -

REACTION SCHEME G

- 25 -

REACTION SCHEME G (cont'd)

5




10

1.  $R_3MgCuCNCl \cdot BF_3$ 2.  $H_2$ , 5% Pd/C

Step K

15




20

1. HCl


2.  $ClCH_2CH_2CH_2COCl$ , pyridine

25



30


NaOH



- 26 -

The oxa isostere compounds of this invention are prepared according to the route outlined in Scheme H. An aminoalcohol 1 is acylated with alpha-chloroacetyl chloride in the presence of trialkylamines to yield amide 2. Subsequent reaction of 2 with a  
5 deprotonation reagent (e.g., sodium hydride or potassium t-butoxide) in an ethereal solvent such as THF provides morpholinone 3. The N-Boc derivative 4 is then obtained by the treatment of 3 with BOC anhydride and DMAP (4-dimethylaminopyridine) in methylene chloride. Alkylation of 4 with  $R^3XL$ , where  $XL$  is a leaving group such as  $Br^-$ ,  $I^-$  or  $Cl^-$  in THF/DME (1,2-dimethoxyethane) in the presence of a suitable  
10 base, preferably NaHMDS [sodium bis(trimethylsilyl)amide], affords 5, which is retreated with NaHMDS followed by either protonation or the addition of an alkyl halide  $R^4X$  to give 6a or 6b, respectively. Alternatively, 6a can be prepared from 4 via an aldol condensation  
15 approach. Namely, deprotonation of 4 with NaHMDS followed by the addition of a carbonyl compound  $R^7R^8CO$  gives the adduct 7. Dehydration of 7 can be effected by mesylation and subsequent elimination catalyzed by DBU (1,8-diazabicyclo[5.4.0]undec-7-ene) or the direct treatment of 7 with phosphorus oxychloride in pyridine to  
20 give olefin 8. Then, catalytic hydrogenation of 8 yields 6a. Direct hydrolysis of 6 with lithium hydrogen peroxide in aqueous THF will produce acid 9b. Sometimes, it is more efficient to carry out this conversion via a 2-step sequence, namely, hydrolysis of 6 in hydrochloric acid to afford 9a, which is then derivatized with BOC-ON  
25 or BOC anhydride to give 9b. The peptide coupling of acid 9b with either an alpha-aminolactone (e.g., homoserine lactone, etc.) or the ester of an amino acid is carried out under the conditions exemplified in the previously described references to yield derivative 10. Treatment of 10  
30 with gaseous hydrogen chloride gives 11, which undergoes acylation with 4-chlorobutyryl chloride. Base treatment leads to lactam formation.



- 27 -

SCHEME H

- 28 -

SCHEME H (CONT'D)



5

9a,  $\text{R}^9 = \text{H}$ 


10

b,  $\text{R}^9 = \text{BOC}$ 

15

10

20


11

25

30

- 29 -

**SCHEME H (CONT'D)**




The thia, oxothia and dioxothia isostere compounds of this invention are prepared in accordance to the route depicted in Scheme I. Aminoalcohol 1 is derivatized with BOC<sub>2</sub>O to give 15. Mesylation of 15 followed by reaction with methyl α-mercaptoacetate in the presence of cesium carbonate gives sulfide 16. Removal of the BOC group in 16 with TFA followed by neutralization with di-isopropylethylamine leads to lactam 17. N-BOC derivative 18 is obtained via the reaction of 17 with BOC anhydride in THF catalyzed by DMAP. Sequential alkylation of 18 with the alkyl halides R<sup>3</sup>X and R<sup>4</sup>X in THF/DME using NaHDMS as the deprotonation reagent produces 19. Hydrolysis of 19 in hydrochloride to yield 20a, which is derivatized with Boc anhydride to yield 20b. The coupling of 20b with an α-aminolactone (e.g., homoserine lactone, etc.) or the ester of an amino acid is carried out under conventional conditions as exemplified in the previously described references to afford 21. Sulfide 21 is readily oxidized to sulfone 22 by the use of MCPBA (m-chloroperoxybenzoic acid). The N-BOC group of either 21 or 22 is readily removed by treatment with gaseous hydrogen chloride. The resultant amine hydrochloride 23 can be converted to the lactam as described above.




- 30 -


## SCHEME I



- 31 -



SCHEME I (CONT'D)

15



25

30



- 32 -

The compounds of this invention inhibit Ras farnesyl transferase which catalyzes the first step in the post-translational processing of Ras and the biosynthesis of functional Ras protein. These compounds are useful as pharmaceutical agents for mammals, especially  
5 for humans. These compounds may be administered to patients for use in the treatment of cancer. Examples of the type of cancer which may be treated with the compounds of this invention include, but are not limited to, colorectal carcinoma, exocrine pancreatic carcinoma, and myeloid leukemias.

10 The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers or diluents, optionally with known adjuvants, such as alum, in a pharmaceutical composition, according to standard pharmaceutical practice. The  
15 compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

For oral use of a chemotherapeutic compound according to this invention, the selected compound may be administered, for  
20 example, in the form of tablets or capsules, or as an aqueous solution or suspension. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch, and lubricating agents, such as magnesium stearate, are commonly added. For oral administration in capsule form, useful diluents include lactose and dried  
25 corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring agents may be added. For intramuscular, intraperitoneal, subcutaneous and intravenous use,  
30 sterile solutions of the active ingredient are usually prepared, and the pH of the solutions should be suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled in order to render the preparation isotonic.

The present invention also encompasses a pharmaceutical composition useful in the treatment of cancer, comprising the

- 33 -

administration of a therapeutically effective amount of the compounds of this invention, with or without pharmaceutically acceptable carriers or diluents. Suitable compositions of this invention include aqueous solutions comprising compounds of this invention and  
5 pharmacologically acceptable carriers, e.g., saline, at a pH level, e.g., 7.4. The solutions may be introduced into a patient's intramuscular blood-stream by local bolus injection.

When a compound according to this invention is administered into a human subject, the daily dosage will normally be  
10 determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for  
15 cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 20 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 10 mg/kg of body weight per day.

20

### EXAMPLES

Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of the invention  
25 and not limitative of the reasonable scope thereof.

The standard workup referred to in the examples refers to solvent extraction and washing the organic solution with 10% citric acid, 10% sodium bicarbonate and brine as appropriate. Solutions were dried over sodium sulfate and evaporated in vacuo on a rotary  
30 evaporator.

- 34 -

EXAMPLE 1Preparation of N-[2(S)-(pyrrolidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine

5

Step A: Preparation of N-4-chlorobutanoylvalyl-isoleucyl-methionine methyl ester

The hydrochloride salt of valyl-isoleucyl-methionine methyl ester was obtained using standard solution phase synthesis methods. To a suspension of this tripeptide (150 mg, 0.38 mmol) in 10 mL methylene chloride at 0°C was added pyridine (61 mL, 0.75 mmol), 43 mL of 4-chlorobutanoyl chloride (0.38 mmol) and 5 drops of DMF. The mixture was stirred for 15 min. at 0°C and then at room temperature overnight. The reaction was worked up in the standard manner to afford 130 mg of crude product. This material was further purified by chromatography on silica gel (98:2 methylene chloride:methanol) and trituration with a mixture of ether and methylene chloride. The solid product weighed 70 mg.

20

Step B: Preparation of N-[2(S)-(pyrrolidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine

The product of Step A was suspended in 5 mL of methanol and 0.6 mL of 1N NaOH was added. After 1 h. the methanol was evaporated, the residue was dissolved in water and filtered. The filtrate was acidified and the product was extracted into ethyl acetate. After standard workup, a solid was obtained, which was washed with water and ether. After drying the solid weighed 25 mg. FAB mass spectrum  $m/z = 430$  (M+1).

30

Anal. Calcd for  $C_{20}H_{35}N_3O_5S \cdot 0.2 H_2O$ :

C, 55.46; H, 8.24; N, 9.70.

Found: C, 55.45; H, 7.88; N, 9.61.

- 35 -

EXAMPLE 2Preparation of N-[2(S)-(piperidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine

5

Step A: Preparation of N-(5-chloropentanoyl)valine methyl ester

Using the method of Example 1, Step A, valine methyl ester and 5-chloropentanoyl chloride were coupled to provide N-(5-chloropentanoyl)valine methyl ester.

10

Step B: Preparation of Methyl 2(S)-(piperidin-2-on-1-yl)-3-methylbutanoate

A solution of 105 mg (0.42 mmol) of the product of Step A in 10 mL of dry THF was treated with 17 mg (0.43 mmol) of 60% NaH in oil dispersion at 0°C for 1 h. TLC analysis indicated starting material was present and 2 mL of DMF was added. After 1.5 h. at 0°C, the solvent was evaporated and the mixture was worked up in the standard to give 90 mg of product.

15

Step C: Preparation of 2(S)-(Piperidin-2-on-1-yl)-3-methylbutanoic acid

The product of Step B was dissolved in 4 mL of methanol and 0.85 mL of 1 N NaOH was added. The mixture was stirred overnight and the solvent was evaporated. The residue was dissolved in water, filtered and acidified. Ethyl acetate extraction and standard workup gave 30 mg of the title compound.

25

Step D: Preparation of N-[2(S)-(piperidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine

30

The product of Step C was coupled to the dipeptide isoleucyl-methionine methyl ester using a standard protocol. Following saponification of the methionine ester, the title compound was obtained. <sup>1</sup>H-NMR of this material indicated that it was a 1:1 mixture of diastereomers, due to racemization of the valine residue.

- 36 -

EXAMPLE 3In vitro inhibition of Ras farnesyl transferase

5 Farnesyl-protein transferase (FTase) from bovine brain  
was chromatographed on DEAE-Sephacel (Pharmacia, 0-0.8 M NaCl  
gradient elution), N-octyl agarose (Sigma, 0-0.6 M NaCl gradient  
elution), and a mono Q HPLC column (Pharmacia, 0-0.3 M NaCl  
10 gradient). Ras-CVLS at 3.5 mM, 0.25 mM [<sup>3</sup>H]FPP, and the indicated  
compounds were incubated with either a partially purified bovine  
enzyme preparation or a recombinant human enzyme preparation. The  
FTase data presented below in Table 1 reflects the ability of the test  
compound to inhibit RAS farnesylation in vitro, as described in  
15 Pompliano, et al., Biochemistry 31, 3800 (1992).

TABLE 1Inhibition of RAS farnesylation by compounds of this invention\*

20	N-[2(S)-(pyrrolidin-2-on-1-yl)-3-methylbutanoyl]- isoleucyl-methionine	2000
25	N-[2(S)-(piperidin-2-on-1-yl)-3-methylbutanoyl]- isoleucyl-methionine	6200


\*(IC<sub>50</sub> is the concentration of the test compound which gives 50%  
inhibition of FTase under the described assay conditions).

30

- 37 -

WHAT IS CLAIMED IS:

1. A compound which inhibits Ras farnesyl-transferase having the formula I:



I

wherein:

V is CH<sub>2</sub>, O, S, HN, or R<sup>7</sup>N;


R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;




- 38 -

X-Y is


5






10



15



20



25

R<sup>7</sup> is

an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

30

Z is

H<sub>2</sub> or O;

m is

0, 1 or 2; and


o is

0, 1, 2 or 3;

or the pharmaceutically acceptable salt thereof.

- 39 -

2. A prodrug of a compound of Claim 1 having the formula II:



II

10

wherein:

V is CH<sub>2</sub>, O, S, HN, or R<sup>7</sup>N;

15







R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;

25

30

- 40 -

X-Y is



R<sub>6</sub> is

a substituted or unsubstituted aliphatic, aromatic or heteroaromatic group such as saturated chains of 1 to 8 carbon atoms, which may be branched or unbranched, wherein the aliphatic substituent may be substituted with an aromatic or heteroaromatic ring;

R<sub>7</sub> is

an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

- 41 -


Z is H<sub>2</sub> or O;

m is 0, 1 or 2; and

o is 0, 1, 2 or 3;

5 or the pharmaceutically acceptable salts thereof.

3. A compound which inhibits Ras farnesyl-transferase having the formula III:



wherein:

V is CH<sub>2</sub>, O, S, HN, or R<sup>7</sup>N;

20







R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;

25

30

- 42 -

X-Y is



R<sup>7</sup> is an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

Z is H<sub>2</sub> or O;

n is 0, 1 or 2;

m is 0, 1 or 2; and

o is 0, 1, 2 or 3;


or the pharmaceutically acceptable salts thereof.

- 43 -

4. A prodrug of a compound of Claim 3 of the formula

IV:

5



10

IV

**wherein:**

15







V is CH<sub>2</sub>, O, S, HN, or R<sup>7</sup>N;

20 R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are independently the side chains of naturally occurring amino acids, including their oxidized forms which may be methionine sulfoxide or methionine sulfone, or in the alternative may be substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, such as allyl, cyclohexyl, phenyl, pyridyl, imidazolyl or saturated chains of 2 to 8 carbon atoms which may be branched or  
25 unbranched, wherein the aliphatic substituents may be substituted with an aromatic or heteroaromatic ring;

30

- 44 -

X-Y is



R7 is

an alkyl group, wherein the alkyl group comprises straight chain or branched chain hydrocarbons of 1 to 6 carbon atoms, which may be substituted with an aromatic or heteroaromatic group;

Z is

H2 or O;

30

n is

0, 1 or 2;

m is

0, 1 or 2; and

o is

0, 1, 2 or 3

or the pharmaceutically acceptable salts thereof.

- 45 -

5. A compound of Claim 1 which inhibits farnesyl-protein transferase which is:

5 N-[2(S)-(pyrrolidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine;  
or

N-[2(S)-(piperidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine;  
10 or the pharmaceutically acceptable salts thereof.

6. A compound which inhibits farnesyl-protein transferase which is:

15 N-[2(S)-(pyrrolidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine  
or the pharmaceutically acceptable salts thereof.

7. A compound which inhibits farnesyl-protein transferase which is:

20 N-([2(S)-(piperidin-2-on-1-yl)-3-methylbutanoyl]-isoleucyl-methionine  
or the pharmaceutically acceptable salt thereof.

8. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 1.

25 9. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 2.

30 10. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 3.



- 46 -

11. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 4.

5 12. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 6.

10 13. A pharmaceutical composition comprising or a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a composition of Claim 7.

15 14. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 8.

20 15. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 9.

16. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 10.

25 17. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 11.

30 18. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

- 47 -

19. A method for inhibiting farnesylation of Ras protein which comprises administering of a mammal in need thereof a therapeutically effective amount of a composition of Claim 13.

5           20. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 8.

10           21. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 9.

15           22. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 10.

20           23. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 11.

          24. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

25           25. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 13.

30

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/10872

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 38/00  
US CL : 514/19; 548/550; 546/216

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/19; 548/550; 546/216

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

USPTO APS

search terms: farnesyl, grangroth, guilhot, desolms, graham, inhibitor#

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 5,238,922 (GRAHAM ET AL.) 24 AUGUST 1993, see entire document.	1-25
A	Science, Volume 260, issued 25 June 1993, G. L. James et al., "Benzodiazepine Peptidomimetics: Potent Inhibitors of Ras Farnesylation in Animal Cells", pages 1937-1942, see entire document.	1-25

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*Z\* document member of the same patent family

Date of the actual completion of the international search

26 OCTOBER 1994

Date of mailing of the international search report

09 JAN 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SHEELA J. HUFF

Telephone No. (703) 308-0196